

BEST AVAILABLE COPY

REMARKS

Claims 29-40 are pending in the application. Claims 1-28 are cancelled, claims 29-36 were previously pending, and new claims 37-40 are added by this amendment. These new claims have support throughout the specification, and specifically at page 4, lines 15-20. No new matter has been added by the addition of these claims.

Formal Matters

The Examiner indicated that the application is not in compliance with the sequence rules because the sequences in Figure 1 are not accompanied by reference to SEQ ID NOS:. The Description of the Drawings section of the application has been amended to include a reference to SEQ ID NOS:1 and 2 in the drawings. In light of this amendment, Applicants request that the Examiner withdraw this objection.

Rejection under 35 U.S.C. § 112, first paragraph

The Examiner rejected claims 29-36 under 35 U.S.C. § 112, first paragraph, as allegedly failing to comply with the enablement requirement. The Examiner contends that claims 29-36 do not enable one skilled in the art to make and/or use the invention.

The Examiner bases the rejection on two premises: (1) that BMP-12, BMP-13, and MP-52 preferentially enhance the growth of tendon and/or ligament tissue; and (2) that neither tendonitis nor carpal tunnel syndrome is characterized by loss of tendon or ligament tissue. The Examiner then argues that tendonitis and carpal tunnel syndrome cannot be treated by enhancing the growth of tendon and/or ligament tissue because the enhancement of tendon/ligament tissue will not reduce the inflammation and pain associated with these conditions. Applicants disagree.

The Examiner's argument that tendonitis and carpal tunnel syndrome may only be treated by agents that directly reduce inflammation and pain is incorrect. Damage to tendon tissue is the overwhelming cause of the inflammatory response and pain that make up the symptoms of tendonitis and carpal tunnel syndrome. The inflammation and pain associated with these conditions does not occur in the absence of tissue damage. In particular, the attached table of definitions describes tendonitis as a tendon strain or tear (Leadbetter, *Clinics in Sports Medicine* 11(2):533, 545 (1992)). Carpal tunnel syndrome is caused by damaged and enlarged tendons and ligaments, which pinch a nerve in the wrist (see Carpal Tunnel Syndrome Information Page from NINDS).

Treating the tendon/ligament damage causing the inflammation and pain will result in an overall reduction in the symptoms of tendonitis and carpal tunnel syndrome. As the Examiner acknowledges, BMP-12, BMP-13, and MP-52 can all enhance tendon and ligament formation. Thus, they can also induce the growth of damaged tendon and ligament tissue. This new tendon/ligament growth will reduce tissue damage and will eventually lead to a reduction in the manifested symptoms of tendonitis and carpal tunnel syndrome, inflammation and pain. Applicants submit that no undue experimentation would be needed to demonstrate that BMP-12, BMP-13, and MP-52 can treat tendonitis and carpal tunnel syndrome, and thus the claims are enabled.

Nevertheless, and in an effort to further prosecution, Applicants have added new claims 37-40, which recite a method for treating a tendon or ligament defect. Applicants respectfully request that the Examiner withdraw the rejection of claims 29-36 under 35 U.S.C. § 112, first paragraph.

Double Patenting Rejection

The Examiner rejected claims 29-36 under the judicially created doctrine of obviousness-type double patenting over claims 1-20 of U.S. Patent No. 5,658,882.

Applicants file herewith a terminal disclaimer under 37 C.F.R. §1.321(c) indicating that the '882 patent and the instant application are commonly owned. Applicants note that any patent issuing from this patent would expire on December 7, 2013, before the expiration date of the '882 patent on August 19, 2014. The terminal disclaimer is submitted solely to confirm that any patent issuing from this application will be enforceable only for and during such period that it and the '882 patent are commonly owned. Applicants respectfully request that, in view of the terminal disclaimer, the rejection of the claims as unpatentable over the '882 patent be withdrawn.

In view of the foregoing amendments and remarks, Applicant respectfully requests reconsideration and reexamination of this application and the timely allowance of the pending claims.

Please grant any extensions of time required to enter this response and charge any additional required fees to our deposit account 06-0916.

Respectfully submitted,

FINNEGAN, HENDERSON, FARABOW,
GARRETT & DUNNER, L.L.P.

Dated:


1/7/05

By:


Elizabeth E. McNamee
Reg. No. 54,696

Attachments:

- Terminal Disclaimer;
- Leadbetter, *Clinics in Sports Medicine* 11(2):533,545 (1992);
- Carpal Tunnel Syndrome Information Page from NINDS.



**CERTIFICATE UNDER 37 CFR § 1.10 OF
MAILING BY "EXPRESS MAIL"**

EV 411565260 US
USPS Express Mail Label Number

January 7, 2005
Date of Deposit

I hereby certify that this correspondence is being deposited with the United States Postal Services "Express Mail Post Office to Addressee" service under 37 CFR § 1.10 on the date indicated above and is addressed to the Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

By: Kathy Meuse
Kathy Meuse

CELL-MATRIX RESPONSE IN TENDON INJURY

Wayne B. Leadbetter, MD

△ NOTICE: THIS MATERIAL MAY BE PROTECTED
BY COPYRIGHT LAW (TITLE 17 U.S. CODE)

The cell-matrix response of tendon to sports-induced trauma is characterized by a broad spectrum of histologic and biochemical adaptations to physical load and use. Historically, *tendinitis* has been the clinical term applied to virtually all painful tendon structures, their synovial sheathes, and adjacent bursae. This tradition is challenged by mounting pathologic evidence that distinguishes between the acute traumatic inflammatory response and the more insidious process of chronic tendon degeneration.^{27, 71, 106, 108} Although this is by no means a resolved issue, the attempt to distinguish between these two processes bears great importance in prescribing appropriate therapeutic intervention and lies at the crux of the difficulties in predicting successful return to sports performance.^{82, 136} There is a significant amount of scientific data on the healing response of acutely traumatized tendon and ligament; however, the literature deals primarily with flexor tendon repaired after laceration in the animal model or human hand. Information on "nontraumatic" injury (i.e., chronic overuse) has been lagging because of the inherent nonoperative nature of the early stages of these conditions and the lack, until recently, of an animal model.^{13, 32} This discussion will focus on tissue events at common sites of clinical treatment, such as the rotator cuff tendon complex of the shoulder, the lateral extensor or medial flexor musculature of the forearm and elbow, the patella tendon, and Achilles and posterior tibial tendon. These sites share common attributes of tendons exposed to repetitive high eccentric loads, a muscle tendon unit that in most cases crosses more than one

From the Department of Orthopaedic Surgery, Georgetown University, Washington, DC; Department of Surgery, Uniformed Services University of the Health Sciences, Bethesda, Maryland; and Shady Grove Center for Sports Medicine and Rehabilitation, Rockville, Maryland

CLINICS IN SPORTS MEDICINE

VOLUME 11 • NUMBER 3 • JULY 1992

533

joint, and some quality of tenuous vascularity.^{2, 19, 33} It is not surprising that such tendons comprise the bulk of treatment requirements and have provided much of the existing biopsy material, such as it is. This article reviews the present understanding of inflammation, repair, and degeneration in tendinopathy. The mechanisms of acute and chronic tendon injury are contrasted and compared. Additionally, the effect of epigenic factors such as age, vascularity, load, use, and rest will be considered. Finally, some theoretic models are provided describing tendon pathophysiology and future therapeutic opportunities. Although some similarities exist in acute trauma,^{12, 15} observations will be restricted to tendon tissue response in contradistinction to the myotendinous junction or tendon insertion.

TENDON INJURY, INFLAMMATION, REPAIR, AND DEGENERATION

Sports "injury" (from the Latin *injure*—to make unjust, not right)¹²⁴ is the loss of cells or extracellular matrix resulting from sports-induced trauma. Injury represents a failure of cell matrix adaptation to load exposure, whether sudden overload or accumulative overload secondary to cyclic overuse.⁸³ "Overuse" and "overload" may not be synonymous terms, because injury can result from excessive and rapid change in use without significant change in resistance—hence, the origin of the term "cumulative trauma disorder," or as I prefer "cumulative cell-matrix adaptive response."⁸ Synovial structures such as the tendon sheath as well as peritenon structures are prone to this form of stress response.^{8, 78, 79} Injuries are divided into acute and chronic patterns according to rate of onset and the mechanism.¹²⁶ *Acute* injuries are typified by a sudden crisis followed by a fairly predictable, although often lengthy, resolution. In tendon, an acute injury often consists of midsubstance ruptures occurring either through aberrant tissue or as the result of high strain rates.^{4, 19} *Chronic* injury is characterized by slow, insidious onset, implying an antecedent subthreshold spectrum of structural damage leading to a crisis episode that is often heralded by pain and/or signs of inflammation. Chronic injury may last months or even years and is distinguished by a persistence of symptoms without resolution. Paratenonitis and tendinitis are typical of such complaints. There appears to be some overlap between acute and chronic injuries, with the bridging stage at 4 to 6 weeks termed the "subacute stage" of injury.⁷³

It is difficult to define sports injury clinically. Furthermore, because tendon injury falls within the domain of an athletic soft-tissue complaint, the athletes' injury is often defined solely by the amount of pain and the inability to perform. Noyes et al¹⁰³ have directed attention to this inadequacy and to the importance of defining the exact anatomic extent and occurrence of tissue injury. Classic signs of inflammation after injury are *not* always present or identifiable.¹³⁶ The complaint is not so much identified with specific structure as within an anatomic area. It is a "painful shoulder," not a painful biceps tendon; it is a

"pai
diate
the j
light
inju
over
reve

stud
dror
vate
play
teno
tend

on fi
of w
a tin
and
form
infla
rege
card
37)
(red:
note
ches
role
is in
accu
and
be a
the s

dam
mat
that
scar
wou
94 Th
as o
"in
integ
rege
reco
tissu
obse
tors

"painful knee," not a painful patella tendon. And although the immediate onus on the examiner is to be competent and accurate in making the physical diagnosis, the elicitation of pain does not necessarily shed light on the exact pathology or mechanism of injury. Hence, tendon injury relating to occult joint instability or dynamic tendon stress overload as may be present with the hyperpronating foot may be revealed only by further analysis.

Rovere's attempt to define tendinitis in a study of theatrical dance students exemplifies this difficulty. He defined tendinitis as "a syndrome of pain and tenderness localized over a tendon, usually aggravated by activities that bring the particular muscle tendon unit into play, usually against resistance. . . . The syndrome is inclusive of tenosynovitis and tenovaginitis as well as actual inflammation of the tendon substance itself."⁸³

"Sports-induced inflammation" (from the Latin *inflammare*, to set on fire)¹²³ is a localized tissue response initiated by injury or destruction of *vascularized* tissues exposed to excessive mechanical load or use. It is a time-dependent evolving process characterized by vascular, chemical, and cellular events leading to tissue repair, regeneration, or scar formation. Clinically observed pathways of sports-induced soft-tissue inflammation include spontaneous resolution, fibroproductive healing, regeneration, or chronic inflammatory response (Fig. 1). The four cardinal signs of acute inflammation were defined by Celsus (AD 14-37) in the often-quoted phrase: "*rubor et tumor cum calore et dolore*" (redness and swelling with heat and pain) (Table 1). It is important to note that this was likely a description of an empyema and fistula of the chest.⁸² Based on historic tradition, pain has assumed a disproportionate role in the definition of inflammation, such that any painful structure is immediately presumed inflamed. It has taken the advent of more accurate noninvasive assessment such as magnetic resonance imaging and the accumulation of surgical biopsy evidence to correct what may be a clinical misinterpretation.^{26, 71, 72, 108} As will be further discussed, the source of connective tissue pain is now known to be multifactorial.⁵¹

Repair of soft-tissue injury has been defined as replacement of damaged or lost cells and extracellular matrices with new cells and matrices.¹³⁸ Regeneration is a form of repair that produces new tissue that is structurally and functionally *identical* to normal tissue. Repair by scar is the postnatal mammalian response to injury, unlike in the fetal wound, which is capable of healing without exuberant scar formation.¹ ⁹⁴ This is especially true in a stable tissue (less than 1.5% mitotic activity) as opposed to a labile tissue such as liver.⁹² Jennings and Hunt⁶³ note "in the post-natal animal, the cellular response to achieve tissue integrity by scar formation after injury overrides the potential for perfect regeneration and return of all function, placing importance on early recovery free of infection; [however] perhaps the ability to regenerate tissue is not lost but is subdued by the necessity for survival." Such observations are the underpinning of the present interest in cell mediators to modulate healing response.




Figure 1. Schema of the theoretical pathways of sports-induced inflammatory response. (From Leadbetter WB: An introduction to sports-induced inflammation. In Leadbetter WB, Buckwalter JB, Gordon SL (eds): Sports-Induced Inflammation: Clinical and Basic Science Concepts. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990; p 2; with permission.)

Degeneration describes a change in tissue from a higher to a lower or less functionally active form.⁸² Such weakened structures are then more vulnerable to sudden dynamic overload or cyclic overloading leading to mechanical fatigue and failure. A prominent source of degeneration is cell atrophy, which is the decrease in the size and/or function of a cell in response to a presence (or lack of) an environmental signal.¹¹⁵ Such down-regulation involves decreased protein synthesis and a decrease in such activities as energy production, replication, storage, and contractility. In sports injury, immobilization is a prominent cause of cell atrophy in tendon.^{41, 45, 114, 140} Additional causes include decreased nutrition, diminished endocrine hormonal influence, persistent inflammation, aging, and denervation (Fig. 2). Reversal of the degenerative process is not a typical feature in degenerative conditions

Table

Her

Rec

Swi

Pai

Los

0

*F

beyond

repre

macr

Pr

Figur

results

is mox

Table 1. RECOGNITION OF THE "CARDINAL SIGNS" OF INFLAMMATION

Heat, <i>Calor</i> —metabolic radiant energy
Redness, <i>Rubor</i> —increased vascularity (angiogenesis) and blood flow
Swelling, <i>Tumor</i> —extracellular edema and matrix changes
Pain, <i>Dolor</i> —stimulation of afferent nerve endings by noxious mediators
Loss of function,* <i>Functi laesa</i> —decreased performance caused by direct damage or inhibiting pain, edema

*First pronounced by Virchow.¹³¹

beyond an undefined cell matrix limit.¹³⁶ Ultimately, degeneration represents a profound imbalance in cell matrix homeostasis.

Chronic inflammation involves the replacement of leukocytes by macrophages, plasma cells, and lymphocytes in a highly vascularized




Figure 2. Cell matrix response to change in functional level. In this model, tendinosis results from a failed cell matrix adaptation to excessive changes in load use. Such failure is modified by both intrinsic and extrinsic factors.

and innervated loose connective-tissue milieu at the site of injury.^{10, 84, 100, 114} Although findings of chronic inflammation are typical in sites such as the lateral epicondylar lesions of the elbow,¹⁰⁰ such responses are not found in all chronic sports injuries.⁷⁰ The mechanism that converts an acute inflammation to a chronic inflammatory process is not known; continued abusive load and irritation may stimulate the local release of cytokines, resulting in both autocrine (cell self-stimulation) and paracrine (stimulation of adjacent cells) modulation of further cell activity.^{3, 10, 29, 34, 47}

Inflammation, degeneration, and repair form a functional spectrum of cell matrix responses with predominance of any one response depending on the mechanism of injury and the homeostatic balance of the tendon tissue.

TENDON INJURY RESPONSE

All sports-related connective tissue injury response occurs in the context of two interrelated categories: (1) macrotraumatic—acute tissue destruction; and (2) microtraumatic—chronic abusive load or use.

In tendon, the mechanism of injury has much to do with the subsequent pathohistologic pattern.

Acute Macrotraumatic Tendon Injury Response

Acute tissue loss or damage results in regeneration, fiber-productive response and repair by scar tissue, or some combination of both.^{56, 63} Factors that contribute to the evolution of chronic inflammation or degenerative change are less understood.⁵⁶ The moment of tissue injury is defined by the onset of vascular disruption and the initiation of the clotting mechanism and platelet activation. A cascade of overlapping processes that has been described as "predictable" includes (1) inflammation, (2) cell replication, (3) angiogenesis, (4) matrix deposition, (5) collagen protein formation, (6) contraction, i.e., remodeling, and, in the case of exposed wounds, (7) epithelization (Fig. 3). In fact, this represents an ideal sequence of events influenced by not only the type of insult but also such facts as age, vascularity, nutrition, genetics, hormonal changes, innervation, and activity level. The literature contains many excellent and exhaustive reviews of the vascular cellular and biochemical events in this process.^{1, 5, 15, 18, 25, 29, 34, 40, 44, 55, 56, 62, 89, 105, 114, 129, 138} Acute connective-tissue injury has been further classified into three phases.^{73, 83, 104:}

Phase 1: Acute vascular-inflammatory response. After wounding, the first reparative "cells" to appear in most vascularized wounds are platelets, which are a prominent source of cell mediators such as platelet-derived growth factor (PDGF), platelet factor 4, insulin-like growth factor 1 (IGF-1), transforming growth factor (TGF- β 1 and β 2), and an uncharacterized chemotractant to endothelial cells.⁵⁷ Activation of the coagulation cascade and formation of a fibrin clot containing fibronectin with crosslinking to collagen is vital to facilitate reparative

Fibrin-

Figur
a vari
is an
relati
hours
regen
musc
Sport

cell
that
Hya.
acts
degr
stim
are
initia
colla
endo
Duri
cell
of ce
cells
may
with
mem
cytol
casca
of pr
react
the
tion.




Figure 3. Ideal wound healing model. Originally derived from the study of skin lacerations, a variety of factors may distort the actual healing sequence in tendon. Although this diagram is an accurate portrayal of cell matrix wound healing events, note that the temporal relationship of the various phases is such that the duration of phase one is measured in hours or a few days, but phase three may extend indefinitely. Normal tendon is not regenerated, however. PMN = polymorphonuclear cell. (Adapted from Gamble JG: The musculoskeletal system: Physiological basics. In Hunter-Griffin L (ed): Athletic Training and Sports Medicine, ed 2. New York, Raven Press, 1988, p 105; with permission.)

cell activity. Fibronectins are a class of noncollagenous glycoproteins that act as adhesive molecules, integrating the extracellular matrix. Hyaluronate, a high-molecular-weight matrix glycosaminoglycan, interacts with fibronectin to create a scaffold for cell migration; later, its degradation by neutrophil hyaluronidases to a smaller molecular form stimulates the angiogenesis that will support fibroblast activity. There are three major consequences of the inflammatory phase: (1) some initial wound strength is provided by crosslinking a fibronectin in collagen; (2) damage tissue from the initial trauma is removed; (3) and endothelial sites and fibroblasts are recruited and stimulated to divide.⁹¹ During this phase, release of complement activates polymorphonuclear cell migration into the extravascular space, providing for the removal of cellular debris and initiating chemotaxis of additional inflammatory cells including the tissue macrophage. In tendon, these phagocytic cells may derive from differentiation of epitenon fibroblasts.⁹⁰ Granules within these leukocytes release hydrolytic enzymes that hydrolyze cell membrane phospholipids, producing arachidonic acid, metabolites, cytokines, proteases, and oxidants (Fig. 4). The resulting *arachidonic acid cascade* is an enzymatically driven sequence, leading to the production of prostaglandins, thromboxanes, leukotrienes, eicosanoids, and slow reacting substance of anaphylaxis (SRS-A)¹¹⁵ (Fig. 5). This cascade is the primary chemical event producing the cardinal signs of inflammation. Initiating in minutes, this phase lasts for essentially as long as the




Figure 4. Mediators of the inflammatory response. (From Fantone JC: Basic concepts in inflammation. In Leadbetter WB, Buckwalter JB, Gordon SL (eds): Sports-Induced Inflammation: Clinical and Basic Science Concepts. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990; with permission.)

body requires to set the stage for repair.⁷ Assuming no coincident infection or repetitive disturbance to the wound, this is usually a matter of 3 to 5 days.

Phase 2: Repair-regeneration. Beginning at 48 hours and lasting up to 6 to 8 weeks, this phase is characterized by the presence of the tissue macrophage, formally a circulating monocyte. This pluripotent cell is the starship of wound repair and is capable of directing essentially the complete sequence of events in this proliferative phase. The macrophage is characteristically mobile, capable of releasing a wide menu of growth factors, chemotactants, and proteolytic enzymes when appropriate or necessary for the activation of fibroblasts and tendon repair. The reparative cell in tendon injury is the tenocyte, which when activated behaves like a modified form of fibroblast or fibrocyte. This cell is the source of collagen production, protein mediators of repair, and matrix proteoglycans. Tenocytes are classified as stable cells, meaning less than 1.5% are mitotically active at any one time.²¹ The cells have a low respiratory quotient and a low rate of collagen turnover.^{28, 47} Bound normally within a dense, linear-oriented collagenous matrix, these cells take on a distinct biologic behavior and a fibroproductive phenotypic expression dictated in part by their deformation and cell shape.⁴ Initially, type III collagen in a woven pattern is rapidly deposited. Type III collagen is characterized by a small fibril that is deficient in crosslinking. The remainder of the repair process is characterized by shift to the deposition of type I collagen, which continues for an indeterminate period in the final maturation phase.⁴⁵ The critical, driving force in this stage of the wounding response is a relative hypoxia in the wound micro environment and rising lactate




Figure 5. Generation of arachidonic acid metabolites. HETEs = hydro-oxyecoaia tetranic acids; HPETEs = hydro-peroxyecoiatetranoic acid compounds; LT = leukotriene. (From Fantone JC: Basic concepts in inflammation. In Leadbetter WB, Buckwalter JB, Gordon SL (eds): Sports-Induced Inflammation: Clinical and Basic Science Concepts. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990; with permission.)

levels contributed in part by the release of large amounts of lactate by the tissue macrophage.⁵⁷

Phase 3: The remodeling maturation phase. This is characterized by a trend toward decreased cellularity and an accompanying decrease in synthetic activity, increased organization of extracellular matrix, and a more normal biochemical profile.⁸⁰ Collagen maturation and functional linear realignment are usually seen by 2 months after injury. In the lacerated flexor tendon, by approximately 4 months after injury there appears to be complete maturation of the repair site, and the fibroblasts revert to quiescent tenocytes. Final biomechanical properties can be reduced by as much as 30% despite this remodeling effort.^{4, 33, 45, 46} Biochemical differences in collagen type and arrangement, water content, DNA content, and glycosaminoglycan content persist indefinitely, and the material properties of these scars never equal those of the intact tendon⁴ (Fig. 6).

Cellular response after tendon laceration may assume two patterns: (1) an extrinsic, callus-like response with proliferation and bridging of the injury by the epitenon cells,^{12, 116} or (2) an intrinsic healing response as the result of endotendon fibroblast repair.^{87, 90} In tendons with a well-developed synovial sheath, wounds producing an accompanying vascular insult such as laceration, crush, avulsion, or rupture will heal by an extrinsic dominant pattern.⁴⁶ This is evidenced by the resulting scar and formation of adhesions.⁶⁶ Excessive fibrosis, its mediation, and its modulation have become the focus of current orthopedic therapy. Passive mobilization and controlled use of load have shown promise in improving tendon and ligament healing.^{125, 134} Studies suggesting the inflammatory response is nonessential and that tendons do possess an intrinsic capacity to heal have direct implications for the biologic potential for cell matrix response in the microtraumatized tendon. Although epitenocyte is actively involved in repair of macrotraumatic lesions, the role of the endotenocyte remains controversial.^{12, 87}

Chronic Microtraumatic Tendon Injury

Microtraumatic tendon injury is distinguished by the observation that degenerative changes are a histologic feature, especially in cases of spontaneous tendon rupture.^{6, 14} This degenerative tendinopathy is thought to be the result of a hypoxic degenerative process involving both tenocyte and matrix components.^{70, 71, 76, 77} Inflammatory cell infiltration and orderly phased wound repair as seen in macrotrauma seem to be absent or aborted. Although much of the histologic evidence for degenerative tendinopathy has been derived from the treatment of spontaneous Achilles tendon rupture,^{6, 26, 27, 30, 71, 77, 78, 81, 108} Kannus et al,⁷¹ in a recently published study of the histopathologic changes preceding spontaneous rupture of tendon in 891 patients (397 Achilles tendons, 302 biceps brachii tendons, 40 extensor pollicis longus tendons, 82 quadriceps tendons and patellar ligaments, and 70 other tendons), found 97% were affected with prior pathologic change. The mean patient age was 49 years. Of interest was the finding of similar pathologic changes in 34% of the otherwise asymptomatic control




Figure 6. Summary of events in macrotraumatic wound response. (From Martinez-Hernandez A: Basic concepts in wound healing. In Leadbetter WB, Buckwalter JB, Gordon SL (eds): Sports-Induced Inflammation: Clinical and Basic Science Concepts. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990, p 78; with permission.)

population. Similar observations by Puddu et al¹⁰⁸ and Clancy²⁷ have led to the definition of a new classification of tendon injury and to the coining of the term "tendinosis": a focal area of intratendinous degeneration that is initially asymptomatic. There has been no convincing evidence for the existence of this form of microtrauma response in the young athlete; however, the profound exposure in youth endurance sports and the prevalence of complaints bear further investigation. Based upon the anatomy of the tendon and its surrounding tissues, it is possible to describe four pathologic conditions (Table 2). This classification emphasizes the distinction between peritenon or synovial inflammation and increasing involvement of the tendons' substance as a likely reflection of failed adaptation to physical load and use, and emphasizes the variable stress responses of tendon structure. Similar pathology has been found in patella tendinosis.^{9, 14, 39}

Blackman et al¹³ have provided recent experimental model evidence in the rabbit for the existence of chronic Achilles paratenonitis with tendinosis, with light microscopic examination revealing degenerative changes in the tendon and increased number of capillaries, infiltrates of inflammatory cells, edema, and fibrosis in the peritenon. Leadbetter⁸⁵ has reported similar findings in the adult athlete with overuse tendon injury requiring surgical treatment. Specimens included Achilles tendon, posterior tibial tendon, digital finger flexor tendon, lateral elbow extensors, medial elbow flexor, patella tendon, and triceps. All specimens displayed varying degrees of the following: (1) tenocyte hyperplasia; (2) a blastlike change in morphology from normal tenocyte appearance; (3) prominent small vessel in-growth with accompanying mesenchymal cells; (4) paravascular collections of histiocytic or macrophagelike cells¹⁸; (5) endothelial hyperplasia and microvascular thrombosis; (6) collagen fiber disorganization with mixed reparation and degenerative change; (7) microtears and collagen fiber separations (Figs. 7 to 12). Inflammatory cell populations were prominent in the synovium and peritendinous structures (Fig. 13), as well as the surrounding areas of intratendinous calcification, and at sites of previous intratendinous steroid injection. Reparative cells were evident in patients with tendinosis and tendinitis pathology despite the coexistent findings of cell matrix degeneration. Polymorphonuclear cells and lymphocytes were minimally present.

The synovial sheath and paratenon is also involved in microtraumatic injury, especially as the result of friction with excitation of the synovial cells. Kvist et al²⁷ studied 16 athletes presenting with peritenodinitis. Increased enzyme activities were mainly found in the fibroblast, inflammatory cells, and vascular walls in the peritenon. The results indicated marked metabolic changes occur with an increased catabolism, lowered pH, and decreased oxygenation of the inflamed areas. Typical findings include fibroexudation with deposition of fibronectin and fibrinogen, proliferation of blood vessels, and in some cases marked endothelial hyperplasia with obliteration of microarterioles. Growth factors have been substantiated to modulate this process^{64-66, 69} (Figs. 14 and 15).

Table 2. TERMINOLOGY OF TENDON INJURY

New	Old	Definition	Histologic Findings	Clinical Signs and Symptoms
Paratenonitis	Tenosynovitis Tenovaginitis Peritendinitis Tendinitis	An inflammation of only the paratenon, either lined by synovium or not Paratenon inflammation associated with intratendinosis degeneration	Inflammatory cells in paratenon or peritendinous areolar tissue Same as I, with loss of tendon collagen fiber disorientation, scattered vascular ingrowth but no prominent intratendinous inflammation Noninflammatory intratendinous collagen degeneration with fiber disorientation, hypocellularity, scattered vascular ingrowth, occasional local necrosis or calcification Three recognized subgroups: each displays variable histology from purely inflammation with acute hemorrhage and tear, to inflammation superimposed upon pre-existing degeneration, to calcification and tendinosis changes in chronic conditions. In chronic stage there may be: 1. Interstitial microinjury 2. Central tendon necrosis 3. Frank partial rupture 4. Acute complete rupture	Cardinal inflammatory signs: swelling, pain, crepitation, local tenderness, warmth, dysfunction Same as I, with often palpable tendon nodule, swelling, and inflammatory signs
Paratenonitis with tendinosis				
Tendinosis	Tendinitis	Intratendinous degeneration due to atrophy (aging, microtrauma, vascular compromise, etc.)		Often palpable tendon nodule that can be asymptomatic, but may also be point tender. Swelling of tendon sheath is absent
Tendinitis	Tendon strain or tear A. Acute (less than 2 weeks) B. Subacute (4-6 weeks) C. Chronic (over 6 weeks)	Symptomatic degeneration of the tendon with vascular disruption and inflammatory repair response		Symptoms are inflammatory and proportional to vascular disruption, hematoma, or atrophy-related cell necrosis. Symptom duration defines each subgroup:

Data from Clancy WG: Tendon trauma and overuse injuries. In Leadbetter WB, Buckwalter JA, Gordon SL, (eds): Sports-Induced Inflammation. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990; and Puddu G, Ippolito E, Postacchini P: A classification of Achilles tendon disease. Am J Sports Med 4:145-150, 1976.




Figure 7. Normal Achilles tendon. Note the linear cell shape and orientation (hematoxylin-eosin, $\times 162.5$).




Figure 8. Achilles tendinosis. Note the tenocyte hyperplasia, cell vacuolation, and blast-like change in morphology characteristic of the stress-responsive tenocyte (i.e., tendon fibroblast) (hematoxylin-eosin, $\times 260$).

Figure
(hemat

Figure
(hemat




Figure 9. Achilles tendinosis. Note prominent small vessel ingrowth and angioneogenesis (hematoxylin-eosin, $\times 162.5$).




Figure 10. Achilles tendinosis. Endothelial hyperplasia and microvascular thrombosis (hematoxylin-eosin, $\times 162.5$).




Figure 11. Achilles tendinosis. Collagen fiber disorganization with coexistent repair and degenerative change (hematoxylin-eosin, $\times 162.5$).

Badalamente et al,^{8,9} in studying the biopsy tissues of typical cumulative trauma disorders including trigger-finger, de Quervain's disease, and carpal tunnel syndrome, identified fibrocartilaginous metaplasia in the trigger-finger and de Quervain's condition but not a




Figure 12. Achilles tendinosis; hyalin degeneration (hematoxylin-eosin, $\times 162.5$).

Figure
inflamm

synov
prese
tissue
tunne

Figure
peritenc




Figure 13. Paratenonitis of a digital flexor tendon of the hand. Note the prominent inflammatory cells with synovial hyperplasia (hematoxylin-eosin, $\times 162.5$).

synovitis. A chondroid metaplasia stress response appeared to be present. Leadbetter found both chondroid metaplasia in the pulley A1 tissue as well as synovitis in the tenosynovium⁸⁵ (see Fig. 13). In carpal tunnel syndrome, a proliferation of the type B secretory synovial cell




Figure 14. Achilles paratenonitis with tendinosis. Note the synovial hyperplasia with peritenon adhesions and tendon swelling.




Figure 15. Clinical appearance of Achilles paratenonitis of the left heel (note tenovagium thickening).

in the tendon sheath was found.⁸ Almekinders et al³ have demonstrated an in vitro capability of the human internal tendon fibroblast to produce inflammatory mediators including prostaglandin E_2 and leukotriene (LTB₄) in response to repetitive motion.

In flat tendons, such as the extensor carpi radialis brevis of the lateral elbow, there are similar findings of intratendinous degeneration with a gray immature edematous and friable scar tissue.¹⁰¹ This tissue is essentially characteristic in appearance of chronic granulation tissues seen in various tendon sites throughout the body. It has the additional characteristic of many small fiber sensory nerves and presumably high concentration of nociceptor stimulators (Fig. 16).

The electronic microscopic appearance of microtraumatic tendon degeneration reveals alterations in the size and shape of mitochondria in the nuclei of the tenocyte. Intracytoplasmic or mitochondrial calcification may be seen.¹³⁵ Dystrophic calcium pyrophosphate salts precipitate in degenerative tendon tissue as a result of mitochondrial injury.¹¹⁵ The resulting calcification is deposited in the collagen matrix as chalky-appearing hydroxypapatite crystals, the "tombstones" of tendon injury. Cytoplasmic vacuoles, lipid deposition, and cell necrosis changes are thought to result from relative hypoxia.⁷¹ Similar changes have been documented in reparative cells, however, after tendon laceration.⁹⁰ Changes in the collagen fibers include longitudinal splitting, disintegra-

Figure 1
of inflam
Dionne F
31:675-6

tion, an
abnorm

Tw
old ma
surgery
demon
organe
swellin
change
aging.¹¹
is aggr
cistern
and pr
produc
degene
metabc
mainte
definec
by pro
with ce




Figure 18. Schema of the positive-feedback relationship that develops during the course of inflammation secondary to sports-related injuries. (From Hargreaves KM, Troullos ES, Dionne RA: Pharmacologic rationale for the treatment of acute pain. *Dent Clin North Am* 31:675-694, 1987.)

tion, angulation with a unique bent-fiber appearance (knicking), and abnormal variation in fiber diameter.^{71, 128}

Two cases from our experience are illustrative. Case 1 is a 42-year-old male jogger found to have an Achilles tendinosis lesion requiring surgery. Biopsy several centimeters proximal to the clinical lesion demonstrated a normal tenocyte linear appearance with fairly normal organelle configurations; however, there is borderline mitochondrial swelling and some variation in collagen fibril size (Fig. 17). The matrix changes seen in this case are also consistent with some quality of aging.^{122, 127} In the same case, at the site of the lesion, fibroblastic activity is aggressive, with pronounced rough endoplasmic reticulum with cisternae dilated with electron-dense material, implying proteoglycan and protein production (Fig. 18). There is pronounced collagen fibril production of variable orientation and size (Fig. 19). Thus, although degeneration has been implicated in tendinosis, some cells still are metabolically active. This suggests a dysfunction of tendon tissue maintenance influenced by environmental factors that are to be further defined. Case 2 (Fig. 20) reveals severe cell stress response as evidenced by prominent lipid storage and even the presence of cholesterol clefts with cell death¹¹⁸ (Fig. 21). It is interesting to note the variability in

Text continued on page 556



Figure 17. Transmission electron micrograph from control biopsy site in Achilles tendon. Note subtle mitochondrial (M) swelling with age-dependent variability in collagen fibril size (V) (osmium, original magnification $\times 38,000$).



Figure 18. Note pronounced swelling and damage (V) (osmium, original magnification $\times 29,250$).




Figure 18. Transmission electron micrograph from tendinosis lesion in Achilles tendon. Note prominent rough endoplasmic reticulum with associated ribosomes and prominent swelling secondary to presumed proteoglycan deposition (ER ↑), mitochondria with cristernal damage (M), and variable collagen fibril production (V) (osmium, original magnification $\times 29,250$).



Figure 19. Transmission electron micrograph of Achilles tendinosis revealing active protein (small arrow) and collagen fibril production (large arrow) (osmium, original magnification $\times 48,750$).

Figure :
lipid stor
consiste
magnific



Figure 20. Transmission electron micrograph of Achilles tendinosis. Note the prominent lipid storage (L), enlarged lysosomal cavities (LY), and swollen disrupted mitochondria (M) consistent with severe cell stress response and impending cell necrosis (osmium, original magnification $\times 10,750$).

structure of the collagen matrix that is found quite far from the clinical lesion (Fig. 22). Because all of the tendon is exposed to the same mode or use pattern mechanically, the implication is that stress adaptations may be occurring throughout the entire course of the tendon structure. The patient becomes symptomatic only at a site where focal overload, hypovascularity, or some other influence further reduces cell-stress response capability. When breakdown exceeds repair, the tendinosis cycle is initiated. It should be noted that this patient had a normal serum lipid profile and no evidence of xanthomatosis.

The pathologic changes in this case were reflected in the appearance of the magnetic resonance scan, but they were remarkably inapparent on gross examination of the tendon at surgery (Figs. 23 to 25).

Acute Versus Chronic Clinical Injury Profile

In addition to the histopathologic pattern of cell response to injury, acute and chronic injuries are distinguished by their clinical injury




Figure 21. Transmission electron micrograph of Achilles tendinosis with cholesterol clefts (C), severe mitochondrial damage (M), myelin forms (MY), and cell necrosis (osmium tetroxide, original magnification $\times 13,750$).

Figure
proxim:
crimp (
implyin
nodule




Figure 22. Transmission electron micrograph of Achilles tendon controlled biopsy site 5 cm proximal to tendinosis lesion. Note variability (V) of collagen fibers despite fairly normal crimp (C) pattern. Prominent endoplasmic reticulum (ER) with evidence of protein production implying significant change can coexist throughout the tendon not only at the tendinosis nodule site (osmium tetroxide, original magnification $\times 8,250$).



Figure 23. Preoperative MR scan of Achilles tendinosis. Note the intratendinous signal.

Figure 24. Achilles 1
fusiform t
peremic t
to intact p




Figure 24. Gross appearance of Achilles tendinosis lesion. Note fusiform tendon swelling and hyperemic tenovagium coincidental to intact plantars tendon.




Figure 25. Achilles tendinosis with intraoperative view of longitudinal internal tenotomy. Note the benign gross appearance.

profiles. The acute injury profile is characterized by a defined time of onset with the trauma episode generally observed as a sudden catastrophic occurrence, such as a collision or contact injury, or in the case of a tendon, a spontaneous midsubstance disruption (Fig. 26). At the moment of injury, pain is likely to be severe. This is typically followed by a period of gradually decreasing pain as inflammation is treated intensively. Pain eventually falls below an arbitrary threshold at which time the patient will feel well. When pain is no longer inhibiting, the athlete will request a return to activity; however, when the biologic curve of wound healing is plotted versus the subjective pain response over time, a potential period of reinjury vulnerability appears. The duration of this period of vulnerability is proportional to the original severity of the structural damage, the rate of healing of the given individual, which is likely to be slower with age, the nature of the target tissue that is injured, and lastly the expected demand or load exposure upon return to sports. The period of vulnerability after an acute injury would in theory be lengthened by any raising of the arbitrary pain threshold of the athlete or by any rapid removal of the subjective pain (for example, with aggressive anti-inflammatory treatment or analgesic treatment). It would be lengthened by the adverse

Pain ↑
Level ↑

(Mo
of li

Figure 2
of an act
tissues s
curved s

effects
by fun
expedit
research
only 70
as muc
can be
absence
muscul

Ch
importa
percept
overex
in pain
disablin
structur
activity
and cu
type of
history
amount
of mild
abusive
transiti
i.e., mi
cation i
first of
injury,




Figure 26. Hypothetical profile of acute macrotraumatic tissue injury. This profile is typical of an acute partial tendon strain or the pattern of healing in other acutely injured connective tissues such as a lateral-collateral ligament sprain in the ankle. Curved dashed line = pain; curved solid line = tissue healing.

effects of any inappropriate immobilization, but it would be shortened by functional rehabilitation or protective bracing that would either expedite fibrogenesis or decrease tensile load on a tendon.⁹⁹ Because research has suggested that an injured connective tissue may attain only 70% to 80% of original structural and biomechanical integrity after as much as 12 months,¹³⁹ the period of vulnerability in these injuries can be lengthy, implying the need for protected activity despite the absence of pain, and an ongoing rehabilitation program to recruit muscular support.¹¹³

Chronic soft-tissue injuries differ from acute injuries in several important ways (Fig. 27). The moment of injury, in the athlete's perception, may be a moment of noxious pain. This often occurs after overexertion such as a long-distance run or intense throwing, resulting in pain becoming insidiously inhibiting over a period, or explosively disabling hours or days after the event. Muscle, tendon, and synovial structures typically evidence this type of stress-response to sports activity. The examiner's inquiry about the preinjury training patterns and cumulative load exposures is critical to understanding why this type of tissue response has been triggered. For instance, a careful history in a marathon distance runner might reveal that an inadequate amount of time was spent in prerace preparation and that several weeks of mild, but not inhibiting, pain symptoms had been generated by abusive training before the actual moment of injury. This is the transitional injury pattern. In theory, subclinical injury and dysfunction, i.e., microtrauma, precede the moment of conscious injury. The implication is that damage has been accumulating for a long time before the first opportunity for medical treatment. This is distinct from acute injury, in which the onset of injury and initial treatment often closely




Figure 27. Profile of chronic microtraumatic soft-tissue injury. This profile is typical of overuse tendon injury. Solid line = percentage of tissue damage.

coincide. The accumulation of repetitive scar adhesions, degenerative change, and adverse effects in chronic microtrauma imply that a recovery will be slower. Again, a period of vulnerability to reinjury results, which is increased when conventional anti-inflammatory measures and reduction pain are applied without regard for the lack of adequate structural integrity. In chronic inflammatory injury, it is the history that provides a proper recommendation and adjustment of activity.⁸²

The Principle of Transition

As previously defined by Leadbetter,⁸²⁻⁸³ the *principle of transition* states that sports injury is most likely to occur when the athlete experiences any change in mode or use of the involved part (Fig. 28). Transitional injury is rate dependent. Sudden ill-timed activity changes, whether in training or during injury recovery, result in an undesired breakdown response and may outstrip tissue morphostatic efforts by imposing overload or overuse demands on the cell matrix environment. There is growing evidence for a cellular disuse transitional response,^{5, 42, 43, 52, 74, 102, 140} as well as an overuse breakdown response.^{16, 30, 49, 54, 62, 84, 107, 141} Colosimo and Bassett³¹ noted that the only factors correlating with the incidence of jumper's knee were hard playing surfaces and an increased frequency of training sessions. Jozsa and associates⁶⁸ have noted that the relationship of complete rupture of the Achilles tendon to a sedentary lifestyle; although the issue is not specifically addressed in their article, it is likely that many of these injuries occurred in the transition from inactivity to activity. Clement et al,³⁰ in a study on Achilles tendinitis and peritendinitis, were able to identify a training




Figure 2
Leadbetter
ed 2. Pe
permissic

error a
these, 1
a return
in an
osseou
of tole
increm
respon
fascial
include
change
differen
in dura
sport t
theory
athletic

Wolf's

Tei
high cc
of pull
of a cy
Achille
movem




Figure 28. Principle of transition: the more rapid the transition, the greater the risk. (From Leadbetter WB: The physiology of tissue repair. In *Athletic Training and Sports Medicine*, ed 2. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1991, p 96; with permission.)

error as the primary etiologic factor in more than 75% of all cases. Of these, the majority represented a sudden increase in mileage. Too rapid a return to activity was also noted as a prime cause of reinjury. Ilizarov,⁵⁸ in an attempt to determine the influence of rate and frequency of osseous distraction on cellular behavior, was able to identify a window of tolerable distraction rate of 1 mm per day with as many as 60 incremental lengthenings, thereby creating a gradual transition stress response. He found the rate effects proportional to the growth of the fascial fibroblast and capillary ingrowth.⁵⁸ Examples of transitional risks include any attempt to increase performance level, improper training, changes in equipment, environmental changes such as new surface or different training altitudes, alterations in frequency, in intensity, and in duration of training, attempts to master new techniques, return to sport too soon after injury, and even body growth itself. Transition theory correlates with current recommendations on periodization in athletic training.¹³⁷

Wolf's Law of Soft Tissue

Tendon structures are subject not only to tensile load but also to high compressive forces. Such compression occurs extrinsically at sites of pulleys and bony prominences⁴; intrinsic compression is the result of a cyclic torque load seen secondary to pronation of the foot in the Achilles tendon, in the anterior cruciate ligament during the rotatory movement of the knee, and in the rotator cuff during torque about the

shoulder. Vogel¹³² has demonstrated an accumulation of large molecular weight proteoglycans in regions of human posterior tibial tendon posterior to the medial malleolus, concluding that there was a synthetic stress response as a physiologic adaptation to compressive forces. Similar findings were noted in the anterior cruciate ligament tissue,¹³² as well as in the bovine flexor tendons, where fibrocartilaginous matrix metaplasia is a common finding.⁷⁵ Woo,¹³⁹ in analyzing the mechanical properties of tendon and ligaments, have theorized an ideal homeostatic level of stress and strain duration to maintain mechanical properties and tendon mass. Frank,⁴² in a review of cellular load response to tendon loading, theorized the existence of a possible degradative pathway blockade by an inhibitor to explain observed proteoglycan accumulations under stress. Quoting Gillard, he noted that the apparent cell-mediated changes in glycosaminoglycan content of different areas of a tendon can be demonstrated in response to both relief and return of compression in tensile loading. Later work verified that this was in proportion to the qualities of the changing load conditions. This has led to further observations that cells, collagen, and matrix in tensile and pressure zones of different tendons are in fact different. Whether these effects are mediated by cytoskeletal deformation or are due to as-yet-unproven local piezoelectric effects of matrix on cells is unknown. It is interesting to note the secondary increases in phospholipase A2 arachidonic acid and prostoglandin E synthesis in loaded bone cells, however, is not known at this mechanism exists in tendon.^{36, 42, 43} Elliott,³⁷ in an extensive review, notes the importance of transmitted tension as a stimulus to the transected tendon; tendon fails to heal if the tendon muscle was excised. Based on these and similar observations, Amadio⁴ has provided a proposed scheme for Wolf's law of soft tissue (Fig. 29).

Epigenic and Genetic Influences on Tendon Cell Matrix Response

There are both epigenetic and genetic influences on tendon cell matrix response. *Epigenic factors* are defined as those factors that can influence the phenotypic expression (i.e., protein production) of the cell without altering the genome.

Vascularity, hormonal influence, and rest may exert an epigenetic influence on tissue injury healing. Aging may be considered to have both an epigenetic and genetic role in injury response.

Aging is best characterized by a failure to maintain homeostasis under conditions of physiologic stress. The salient characteristic of aging is not so much a decrease in basal functional capacity as it is in a reduced ability to adapt to environmental stress.^{114, 115}

Menard and Stanish⁹⁵ note that the tendon collagen fiber possesses all of the cross linkages it will ever have shortly after its synthesis.¹¹ During maturation, reducible cross linkages gradually stabilize.³⁸ This results in a less compliant collagen fiber subject to shear stress injury. Aging also is accompanied by significant decrease in tendon glycosaminoglycan concentration. There are extensive data on the aging of

Figure
and liga
and Clir

cells in
decrea
preser
matrix
not pr
by ph
muno.
increa
overall
corres
glycar
fibers
and r
regula
howev
may c
old ac
norma
there
more
conter
V
tendo.
of the
pressi
tershe
an ex
and lo




Figure 29. Theoretical schema for Wolff's law of soft tissue. (From Amadio PC: Tendon and ligament. In Cohen IK, Diegelmann RF, Lindblad WJ (eds): Wound Healing: Biochemical and Clinical Aspects. Philadelphia, WB Saunders, 1992, p 388.)

cells in vitro.^{50, 53, 88, 96, 98, 112} Collagen synthesis has long been thought to decrease with age; however, this decrease may be overcome in the presence of ascorbic acid. Generally, aging results in changes in the matrix integrity and in the rate of wound healing, although aging does not prevent adequate clinical wound healing, which can be stimulated by physical training. Ippolito^{59, 60} has documented morphologic, immunologic, and biochemical aging changes. With aging, collagen fibers increase in diameter, vary more in thickness, and there may be an overall increase in insoluble collagen.^{59, 60} These morphologic changes correspond to biochemical changes that include a decrease in proteoglycans and a decrease in water content. Parallel changes in elastic fibers also occur. With age, adaptation requires a longer interval of rest and recovery. This is presumably related to the documented down regulation in the cellular biology of the older athlete. With injury, however, the biochemical and morphologic character of tendon tissue may change. In comparing an area of tendon degeneration in a 25-year-old adult with spontaneous Achilles tendon rupture with that of a normal 24-year-old, Ippolito^{59, 60} noted that in the area of tendinosis there was a 34% loss of collagen and an increase in proteoglycans of more than 100% with a significant increase in water and glycoprotein content.

Vascularity has long been thought to play a prominent role in tendon degeneration,^{35, 79, 121, 129} especially in the supraspinatus portion of the rotator cuff, in the Achilles tendon, and at sites of extrinsic bone pressure. Since the injection study of Rathbun and McNab,¹⁰⁹ a watershed area in the distal supraspinatus tendon had been offered for an explanation of the etiology of rotator cuff degeneration. Chanski and Ionnotti,²³ in an extensive review, conclude that the vascularity in

the critical zone of the supraspinatus tendon is actually hypervascular secondary to a low-grade inflammatory incitation with neovascularization after mechanical irritation.²³ Brooks et al¹⁷ likewise came to the conclusion that no significant difference existed between the vascularity of the supraspinatus portion of the rotator cuff, and that factors other than vascularity were important in the pathogenesis of supraspinatus tendon rupture. These assertions tend to shed a different light on the theory of hypoxic intratendinous degeneration and the etiology of tendinosis. Focal load influences on cell matrix metabolism may play as great a role as any proposed diminished vascularity. Wilson and Goodship¹³⁸ have measured a core temperature increase of 5°C to 9°C secondary to hysteresis energy losses in the equine superficial digital flexor tendon during exercise. This radiant energy equals 10% of the lost elastic energy upon unloading and is theorized to be potentially cytotoxic. Further research is needed to resolve this question.¹⁸ Whereas there seems to be a preponderance of support for a diminished microvascular debt in the central core of the round tendon and in the distal third of the Achilles tendon,¹¹⁹ controversy has arisen as to the role of hypovascularity in rotator cuff and supraspinatus tendon degeneration¹²⁹ (Figs. 30 and 31).

Hormonal influence on tendon biology primarily relates to estrogen and to insulin. It has been suggested that diminished estrogen levels, premature menopause, or premenopausal hysterectomy may be associated with the incidence of tendinosis in women; no other data are



Figure 30. Intra-operative view of Achilles tendinosis linear internal tenotomy. Note that perfusion from the internal tendon vasculature is hypovascular relative to the epitendon.

Figure :
Note the
observa
Scand 1

availat
to heal
Re
tendor
of the
Althou
repara
are pr
the te
betwe
includ
classifi
recove
of freq
injury.
athlete
ified lo
any su
Gr
matrix
called
failed
15% o
overus
alis br
drome
Querv




Figure 31. Cross-sectional injection of histologic study of an Achilles tendon vasculature. Note the peripheral distribution with less internal perfusion. (From Schatzker J: Intravital observation on the microvascular anatomy and microcirculation of the tendon. *Acta Orthop Scand* 126(suppl):1-23, 1969; with permission.)

available to support this contention.¹⁰⁰ In addition, diabetics are known to heal with some difficulty.⁸⁹

Rest has long been clinically recognized to aid the patient with a tendon injury. The beneficial role of rest in the therapeutic intervention of the inflammation repair process is experimental and undefined. Although it may be said that rest does not heal; theoretically, cell reparative efforts may catch up in the face of rest. The effects of rest are probably multifactorial and may include improved vascularity in the tendon at rest or represent an improved morphostatic balance between matrix degradation and production. Different forms of rest include total abstinence, protected activity, or altered activity. Such classifications imply a modulation in cell matrix load signal and load recovery phase. There is evidence that repetitive motion and variation of frequency (i.e., cycles) may create a positive reparative signal post-injury.⁴⁶ Absolute rest or abstinence does not de facto increase the athlete's potential to tolerate renewed load during participation. Modified load rehabilitative prescription has been shown to be important to any successful return to sports performance.^{30, 33, 113, 126}

Genetic influences are implicated in the modulation of tendon cell matrix response based primarily on clinical observations. Nirschl¹⁰⁰ has called attention to the mesenchymal syndrome as a potential cause of failed healing. Tendinosis appears in multiple sites in approximately 15% of patients and in sites not necessarily subjected to obvious overuse. An association among lateral epichondral extensor carpi radialis brevis tendinosis, rotator cuff degeneration, carpal tunnel syndrome, cervical and lumbar disk degeneration, plantar fasciosis, de Quervain's syndrome, and trigger-finger tendinosis has been postu-

lated. Jozsa et al⁶⁷ have found blood type O to be statistically related to tendon rupture. It is interesting, as noted by Singer and Jones,¹²⁰ that Achilles tendon rupture in children is uncommon and has been encountered only in children whose parents had experienced tendon rupture. Young adult herniated disk syndrome is often seen in the presence of a familial history (author's personal observation). An underlying collagen diathesis has been theorized. In the author's experience, the significance of the mesenchymal theory is in recognizing the patient who presents with frequent tendon complaints disproportionate to the level of activity. In addition to ruling out systemic disease, these patients are unusually vulnerable and must be counseled as to proper participation and moderation in their activity.

The factors leading to potential failed healing response are both intrinsic and extrinsic and are summarized in Table 3.

Sources of Pain in Tendon Injury

The sources of tendon pain are multifactorial and include both paratenon as well as intrinsic cellular and biomechanical deformation sources. The synovial site is capable of secreting a wide array of inflammatory mediators, in particular interleukin-1 and prostaglandin E₂. How movement and friction activate these cells is not known,^{41, 44, 61, 77, 111} but particles of cartilage and the degradation of other protein molecules may incite a synovial reaction.¹¹¹ The intrinsic tendon fibroblast is capable of producing inflammatory mediator proteins under repetitive stress.³ Elongation underload of the tendon beyond its elastic limit may trigger nociceptors as well as myotendinous reflexes.^{24, 33} There would appear to be a distinction between the pain due to inflammation and that due to degeneration (author's unpublished

Table 3. FACTORS LEADING TO FAILED TENDON HEALING

Intrinsic	
	Vascular vulnerability
	Limited cell function potential
	Limited cellularity
	Aging
	Genetic predisposition (mesenchymal syndrome, collagen crosslinking, etc.)
	Irreversible change, cystic calcific degeneration
	Hormonal
	Other (autoimmunity, etc.)
Extrinsic	
Overt	
	Continued self-abuse
	Improper training (overstimulus or inadequate stimulus)
	Improper technique
	Improper equipment
	Harsh environment
Covert	
	Joint instability
	Extrinsic pressure
	Biomechanical fault

Abnormi
Motion
Force
Vibrat
Fatigu

Figure
sympos
Back P
with per

observ
with c
mator
genera
matrix
deform
A
injury

Furthe

A
regard
previo
long-t
types
matter
qualifi
wound
initial
synovi




Figure 32. Possible mechanisms of musculoskeletal pain. (From White AA III: The 1980 symposium and beyond. In Frymoyer JW, Gordon SW (eds): *New Perspectives in Low Back Pain*. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1989, pp 3-17; with permission.)

observation). The pain of inflammation evolves from excessive loading with cell matrix and distal damage producing the onset of the inflammatory cascade typical of macrotraumatic or synovial irritation.¹³³ Degenerative pain evolves from excessive cyclic loading, which results in matrix molecular damage, loss of tissue strength, resultant increased deformation with loading, and stimulation of mechanoreceptors.

A scheme for musculoskeletal pain that is applicable to tendon injury has been provided (Fig. 32).

Further Clinical Implications

A statement that tendon healing progresses in a typical pattern regardless of the mechanism of injury²⁴ is untenable in light of the previous discussion. The statement that the body's immediate and long-term responses to physical trauma are essentially the same for all types of athletic injuries,¹⁵ or that a tissue's response to injury, no matter what the cause or type of injury, is inflammatory,⁵⁴ must be qualified. The type of injury has a great deal of influence over the wound healing process.⁶³ The size of the wound, the extent of the initial tissue damage, the presence or absence of any coincident contaminovial structure versus extra-articular tendon, may dictate the rate of




Figure 33. Hypothetical model of tendon lesions. A, Intratendinous shear lesion. B, Pinhole tear. C, Gap. D, Intratendinous cystic degeneration (tendinosis). Different mechanisms of healing repair response are implied.

recovery and its adequacy (Fig. 33). The initial events in the resolution of microtraumatic wounds, in particular the metabolic activity and pattern of protein synthesis of the mesenchymal cell population found in tendinosis lesions, is not well understood.

The tenocyte has been theorized to have limited capability for mitosis or increased collagen production beyond maintenance and has been called an end-stage cell.²⁸ Aging tenocytes show a marked decrease




Figure 34. The role of TGF- β in the stimulation of extracellular matrix. (From Sporn MB, Roberts AB: Transforming growth factor- β : Multiple actions and potential clinical applications. JAMA 262:938-941, 1989; with permission of the American Medical Association, Inc.)

In

in intr
cence
in fib
howev
tive re
or afte
increa
gled v
ization
tenocy
inhere
traum
therap
A
is not




Figure 35: Theoretical application of growth factor therapy in tendon injury.

in intracytoplasmic organelles responsible for protein synthesis. Senescence in the tenocyte parallels the known aging phenomenon occurring in fibroblasts throughout the body.²⁰ Other histopathologic studies, however, including our own,⁸⁵ have suggested that significant reparative response capability resides within the activated tenocyte postinjury or after overload/overuse stimulation. The findings of hypercellularity, increased organelle activity, and patterns of scar deposition intermingled with scattered cell necrosis, degeneration, and collagen disorganization suggest a different theory, namely that it is the imbalance of tenocyte reparative efforts and a failed adaptation as opposed to an inherent lack of healing capability that limits the outcome in cumulative traumatic injury. If such capability still resides within the genome, then therapeutic efforts can target its potentiation.

As with other connective tissues, the metabolism of tendon tissue is not regulated by a dominant central control mechanism.²¹ Rather,

tendon cells seem to function in an interdependent way with their matrix. To what degree mutual local control over neighboring cells exists is a matter for ongoing investigation. Normally, a homeostatic balance is maintained between matrix synthesis and degradation; in stressed states, an adaptive response is elicited. Cytokines (which may be defined as soluble products, mostly protein peptides, released from one cell that can modulate the activity of other cells [paracrine] or the same cell [autocrine]) play a critical role in the process of cell communication. Individual cytokines may have multiple biologic activities, and several cytokines may share common functional properties. Cytokines are best understood as analogous to letters in an alphabet, with cellular messages changing according to sequences and the interaction between several cytokines simultaneously. In the open system of connective tissue metabolism, clinical attempts to modulate wound healing have been challenging. The same peptide may affect a cell entirely differently in vitro versus in vivo. Based upon these observations and the observations of Sporn and Roberts and others,^{97, 110, 124} a theoretic model can be proposed for the pathogenesis of sports-induced tendinosis as well as the potential therapeutic role of growth factors, notably transforming growth factor beta (Figs. 34 and 35).

CONCLUSION

The major mysteries of injury and repair have always been and continue to be *what* in the injury stimulates repair, and *how* does the wound recognize that it is no longer needed—that is, *why* does healing stop?⁵⁶ Once thought to be inert biologically, tendons are best appreciated as a heterogeneous group of structures with variations in cell character, collagen orientation, collagen cross linking, vascularity, configuration, load pattern, biomechanical profile, shape, and the presence or absence of a synovial lining. Tendons have the *potential* to manifest one or several stress adaptation behaviors depending upon the mechanism of injury and the nature of the continuing sports demand. Transition plays an important role in the determination of this process.

Since Vichow's work,¹³¹ modern pathology has defined injury in cellular terms. The study of tendon injury is the study of cell injury and its matrix milieu as an expression of pre-existing capacity to adapt to such injury on the part of the injured or intact cell.¹¹⁴ Today, sports injury is further defined by the degree to which normal mechanical properties in function are lost as a consequence of both cell and matrix injury. The modulation of these cell matrix responses regardless of the method provides an intriguing challenge.

ACKNOWLEDGMENT

The author would like to thank Ingrid Major, Janelle Owen, and K. Blanch Emore for their excellent secretarial assistance.

References

1. Adz
2. Allr
Car
p 15
3. Alir
rep
4. Am
(eds
ders
5. Arn
Mo
6. Arn
case
7. Ass
trar
US
8. Bad
cun
dise
9. Bad
ten
10. Bail
dev
390
11. Bail
of c
12. Ban
met
and
13. Bla
osis
14. Bla
4:66
15. Boc
Mir
16. Bro
Ort
17. Bro
vas
18. Bry
19. But
In I
Ins
20. But
age
ma
Ort
21. Cas
anc
Fet
22. Cel
rab
23. Ch
10:
24. Ch
14:

References

1. Adzick NS, Longaker MT: Fetal Wound Healing. New York, Elsevier, 1992
2. Allman F: Impingement, biceps, and rotator cuff lesions. In Zarins B, Andrews JR, Carson WG (eds): Injuries to the Throwing Arm. Philadelphia, WB Saunders, 1985, p 158
3. Almekinders LC, Banes AJ, Ballinger CA: Inflammatory response of fibroblasts to repetitive motion. *Trans Orthop Res Soc* 17:678, 1992
4. Amadio PC: Tendon and ligament. In Cohen IK, Diegelmann RF, Lindblad WJ (eds): Wound Healing: Biochemical and Clinical Aspects. Philadelphia, WB Saunders, 1992, p 384
5. Arnheim DD: Modern Principles of Athletic Training. St. Louis, Times Mirror/Mosby, 1989, p 230
6. Arner O, Lindholm A: Subcutaneous rupture of the Achilles tendon: A study of 92 cases. *Acta Chir Scand Suppl* 239(suppl):1-51, 1959
7. Assoian K, Fleurdelys BE, Stevenson HC, et al: Expression of secretion of type B transforming growth factor by activated human macrophages. *Proc Natl Acad Sci USA* 84:6020-6024, 1987
8. Badalamente MA, Sampson SP, Dowd A, et al: The cellular pathobiology of cumulative trauma disorders/entrapment syndromes: Trigger finger, DeQuervain's disease and carpal tunnel syndrome. *Trans Orthop Res Soc* 17:677, 1992
9. Badalamente MA, Sampson SP, Hurst LC, et al: The pathobiology of stenosing tenosynovitis of the human A1 pulley. *Trans Orthop Res Soc* 14:199, 1989
10. Bailey AJ, Bazin S, Delaunay A: Changes in the nature of the collagen during development and resorption of granulation tissue. *Biochim Biophys Acta* 328:383-390, 1973
11. Bailey AJ, Robins SP, Balian G: Biological significance of the intermolecular crosslinks of collagen. *Nature* 251:105-109, 1974
12. Banes AJ, Donlon K, Link GW, et al: Cell populations of tendon: A simplified method for isolation of synovial cells and internal fibroblasts. Confirmation of origin and biologic properties. *J Orthop Res* 6:83-94, 1988
13. Blackman C, Boquist L, Friden J, et al: Chronic Achilles paratenonitis with tendinosis: An experimental model in the rabbit. *J Orthop Res* 8:541-547, 1990
14. Blazina ME, Kerlan RK, Jobe FW, et al: Jumper's knee. *Orthop Clin North Am* 4:665, 1973
15. Booher JM, Thibodeau GA: Athletic Injury Assessment, ed 2. St. Louis, Times Mirror/Mosby, 1989, p 114
16. Brody DM: Techniques in the evaluation and treatment of the injured runner. *Orthop Clin North Am* 13:541-558, 1982
17. Brooks CH, Revell WJ, Heatley FW: A quantitative histological study of the vascularity of the rotator cuff tendon. *J Bone Joint Surg* 74B:151-153, 1992
18. Bryant WM: Wound healing. *CIBA Clinical Symposium* 29:1-36, 1977
19. Butler DL, Groodes ES, Noyes FR, et al: Biomechanics of ligaments and tendons. In Hutton RS (ed): Exercise and Sports Sciences Reviews, Washington, DC, Franklin Institute Press, 1978, pp 125-181
20. Butler DL, Siegel A: Alterations in tissue response: Conditioning effects at different ages. In Leadbetter WB, Buckwalter JA, Gordon SL (eds): Sports-Induced Inflammation: Clinical and Basic Science Concepts. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990, pp 713-730
21. Castor CW: Regulation of connective tissue metabolism. In McCarty DJ (ed): Arthritis and Allied Conditions: A Textbook of Rheumatology, ed 11. Philadelphia, Lea & Febiger, 1989, pp 256-272
22. Cetta G, Tenn R, Zanabon G, et al: Biochemical and morphological modifications in rabbit Achilles tendon during maturation and aging. *Biochem J* 204:61-67, 1982
23. Chansky HA, Iannotti JP: The vascularity of the rotator cuff. *Clin Sports Med* 10:807-822, 1991
24. Chaytor R, Stanish WD: Clinical treatment of tendinitis. *Sports Medicine Digest* 14:2, 1992

25. Christie AL: The tissue injury cycle and new advances. Toward its management in open wounds. *Journal of the National Athletic Trainers Association* 26:274-278, 1991
26. Clancy WG, Neidhart D, Brand RL: Achilles tendinitis in runners: A report of five cases. *Am J Sports Med* 4(2):46-57, 1976
27. Clancy WG: Tendinitis and plantar fascitis in runners. In D'Ambrosia R, Drez D: *Prevention and Treatments of Running Injuries*. Thorofare, NJ, Charles B. Slack, 1982, p 77
28. Clancy WG: Tendon trauma and overuse injuries. In Leadbetter WB, Buckwalter JA, Gordon SL (eds): *Sports-Induced Inflammation: Clinical and Basic Science Concepts*. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990, p 609
29. Clark RAF, Henson PM (eds): *The Molecular and Cellular Biology of Wound Repair*. New York, Plenum Press, 1988
30. Clement DB, Taunton JE, Smart GW: Achilles tendinitis and peritendinitis: Etiology and treatment. *Am J Sports Med* 12:179, 1984
31. Colosimo AJ, Bassett III, FH: Jumper's knee: Diagnosis and treatment. *Orthop Rev* 19:139-149, 1990
32. Curwin S: Models for use in studying sports-induced soft tissue inflammation: In Leadbetter WB, Buckwalter JA, Gordon SL (eds): *Sports-Induced Inflammation: Clinical and Basic Science Concepts* Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990, pp 103-121
33. Curwin S, Stanish W: *Tendinitis: Its Etiology and Treatment*. Lexington, MA, Collamore Press, 1984
34. DiSabatino CA Jr: Inflammation. In Albrint JA, Brand RA (eds): *The Scientific Basis of Orthopaedics*. New York, Appleton-Century-Crofts, 1979, pp 415-436
35. Edwards DAW: The blood supply and lymphatic drainage of tendons. *Anatomy* 80:147-152, 1946
36. Edwards P, Mirriari A, Mayas J, et al: Mechanical loading alters ligament cell shapes. In *Trans Combined Meetings Orthop Res Soc USA, Japan and Canada*, October 21-23, 1991, p 57
37. Elliott DH: Structure and function of mammalian tendon. *Biological Review* 40:392-421, 1965
38. Eyre DR, Paz MA, Gallop PM: Cross-linking in collagen and elastin. *Annu Rev Biochem* 53:717-748, 1984
39. Ferretti A, Ippolito E, Mariani P, et al: Jumper's knee. *Am J Sports Med* 11:58-62, 1983
40. Flynn JE, Graham JH: Healing of tendon wounds. *Am J Surg* 109:315-324, 1965
41. Fox RJ, Lotz M, Carson DA: Structures and function of synoviocytes. In McCarty DJ (ed): *Arthritis and Allied Conditions: A Textbook of Rheumatology*, ed 11. Philadelphia, Lea & Febiger, 1989, pp 273-287
42. Frank CB, Hart DA: Cellular response to loading. In Leadbetter WB, Buckwalter JA, Gordon SL (eds): *Sports-Induced Inflammation: Clinical and Basic Science Concepts*. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990, p 555
43. Frank C, MacFarlane B, Edwards P, et al: A quantitative analysis of matrix alignment in ligament scars: A comparison of movement versus immobilization in an immature rabbit model. *J Ortho Res* 9:219-227, 1991
44. Gallin JJ, Goldstein IM, Snyderman R (eds): *Inflammation: Basic Principles and Clinical Correlates*. New York, Raven Press, 1988
45. Gamble JG: *The Musculoskeletal System: Physiological Basics*. New York, Raven Press, 1988
46. Gelberman R, Goldberg V, An KN, et al: Tendon. In Woo SL-Y, Buckwalter JA (eds): *Injury and Repair of the Musculoskeletal Soft Tissue*. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1988, p 5
47. Gerber G, Gerberg G, Altman KI: Studies on the metabolism of tissue proteins: I. Turnover of collagen labeled with poline-U-C14 in young rats. *J Biol Chem* 235:3653-2656, 1960
48. Goldring MB, Goldring SR: Skeletal tissue response to cytokines. *Clin Orthop Rel Res* 258:245, 1990
49. Guten GR, Pietrocarlo T, Julson J: Change, alignment, twisting and speed, foot

orth
1991
50. Hall
67
51. Har
Lea
Clin
paet
52. Har
and
16:1
53. Hay
54. Her
239,
55. Hol
dian
16:6
56. Hur
App
57. Hur
Linc
58. Iliza
infl
1985
59. Ipp
bioc
62A
60. Ipp
of t
Trai
61. Ipp
case
62. Jam
1978
63. Jenu
Log
64. Joyc
folk
Trai
65. Joyc
B2,
Soc
66. Joyc
folk
Trai
67. Jozs
tenc
68. Jozs
tenc
J Sp
69. Jozs
Ortl
70. Jozs
Ortl
71. Kan
tenc
72. Kee
rup
73. Kell
18:4

- orthotics for knee pain. American Medical Athletic Association Quarterly 6(1):Oct 1991, p 6
50. Hall DA: The Aging of Connective Tissue. London, Academic Press, 1976, pp 13-67
 51. Hargreaves KM: Mechanisms of pain sensation resulting from inflammation. In Leadbetter WB, Buckwalter JA, Gordon SL (eds): Sports-Induced Inflammation: Clinical and Basic Science Concepts. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990, p 383
 52. Harper J, Amiel D, Harper E: Changes in collagenase and inhibitor in ligaments and tendon during early development of stress deprivation. Trans Orthop Res Soc 16:114, 1991
 53. Hayflick L: Cell aging. Annual Review of Gerontology and Geriatrics 1:26-67, 1980
 54. Herring SA, Nilson KL: Introduction to overuse injuries. Clin Sports Med 6:225-239, 1987
 55. Holmes G, Sharkley N, Syftestad G, et al: Changes in collagen crosslinks, febril diameter and mechanical strength in aging chicken tendons. Trans Orthop Res Soc 16:620, 1991
 56. Hunt TK, Dunphy JE (eds): Fundamentals of Wound Management. New York, Appleton-Century-Crofts, 1979
 57. Hunt TK, Hussain Z: Wound micro environment. In Cohen IK, Diegelmann RF, Lindblad WJ (eds): Wound Healing Philadelphia, WB Saunders, 1992, p 274
 58. Ilizarov GA: The tension-stress effect on the genesis and growth of tissues: II. The influence of the rate and frequency of distraction. Clin Orthop Rel Res 239:263-285, 1989
 59. Ippolito E, Natali PG, Postacchini F, et al: Morphological, immunological, and biochemical study of rabbit Achilles tendon at various ages. J Bone Joint Surg 62A:583-598, 1980
 60. Ippolito E, Postacchini F, Ricciardi-Pollini PT: Biomechanical variations in the matrix of human tendons in relation to age and pathological conditions. Ital J Orthop Traumatol 1:133-139, 1975
 61. Ippolito E, Ricciardi-Pollini PT: Invasive retrocalcaneal bursitis: A report of three cases. Foot Ankle 4:204-228, 1984
 62. James SL, Bates BT, Ostering LR: Injuries to runners. Am J Sports Med 6:40-50, 1978
 63. Jennings RW, Hunt TK: Overview of post-natal wound healing. In Adzick NS, Logaker MT (eds): Fetal Wound Healing. New York, Elsevier, 1992, p 25
 64. Joyce ME, Pruitt DC, Heydemann A, et al: Growth factor synthesis by epitenocytes following flexor tendon injury: A proposed mechanism for adhesions formation. Trans Orthop Res Soc 16:109, 1991
 65. Joyce ME, Pruitt DC, Manske PR: Synthesis and in situ localization of TGF-B1, TGF-B2, PDGF-A, PDGF-B, and SFGF during flexor tendon healing. Trans Orthop Res Soc 17:680, 1992
 66. Joyce ME, Pruitt DL, Russell J, et al: Growth factor synthesis by epitenocytes following flexor tendons injury: A proposed mechanism for adhesions formation. Trans Orthop Res Soc 16:109, 1991
 67. Jozsa L, Balint JB, Kannus P, et al: Distribution of blood groups in patients with tendon rupture. J Bone Joint Surg 71B:272, 1989
 68. Jozsa L, Kvist M, Balint BJ, et al: The role of recreational sport activity in Achilles tendon rupture. A clinical, pathoanatomical and sociological study of 292 cases. Am J Sports Med 17:338, 1989
 69. Jozsa L, Lehto M, Kannus P, et al: Fibronectin and laminin in Achilles tendon. Acta Orthop Scand 60:469, 1989
 70. Jozsa L, Reffy A, Kannus P, et al: Pathological alterations in human tendons. Arch Orthop Trauma Surg 110:15, 1990
 71. Kannus P, Jozsa L: Histopathological changes preceding spontaneous rupture of a tendon. J Bone Joint Surg 73A:1507-1525, 1991
 72. Keene JS, Lash EG, Fisher DR, et al: Magnetic resonance imaging of Achilles tendon ruptures. Am J Sports Med 17:333-337, 1991
 73. Kellert J: Acute soft tissue injuries—a review of the literature. Med Sci Sports Exerc 18:489-500, 1986

74. Klein L, Player JS, Heiple KG, et al: Isotopic evidence for resorption of soft tissues and bone in immobilized dogs. *J Bone Joint Surg* 64A:225-230, 1982
75. Koob TJ, Vogel KG, Thurmond FA: Compression loading in vitro regulates proteoglycan synthesis by fibrocartilage in tendon. *Trans Orthop Res Soc* 16:49, 1991
76. Koob TJ, Vogel KG, Thurmond FA: Compression loading in vitro regulates proteoglycan synthesis by fibrocartilage in tendon. *Trans Orthop Res Soc* 165:49, 1991
77. Kvist M, Jozsa L, Jarvinen MJ, et al: Chronic Achilles paratenonitis in athletes: A histological and histochemical study. *Pathology* 19:1-11, 1987
78. Kvist MH, Lehto MUK, Jozsa L, et al: Chronic Achilles paratendinitis. *Am J Sports Med* 16:616-623, 1988
79. Lagergren C, Lindholm A: Vascular Distribution in Achilles tendon—angiographic and microangiographic study. *Acta Chir Scand* 116:491-495, 1958
80. Laurent TC: Structure, function and turnover of the extracellular matrix. *Adv Microcirc* 13:15-34, 1987
81. Leach RE, James S, Wasilewski S: Achilles tendinitis. *Am J Sports Med* 9:93, 1981
82. Leadbetter WB: An introduction to sports-induced inflammation. In Leadbetter WB, Buckwalter JA, Gordon SL (eds): *Sports-Induced Inflammation: Clinical and Basic Science Concepts*. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990, pp 3-23
83. Leadbetter WB: Physiology of tissue repair. In *Athletic Training and Sports Medicine*, ed 2. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1991, p 96
84. Leadbetter WB: Getting ahead of the injury. *Emergency Medicine* 15:7-39, 1979
85. Leadbetter WB: The pathohistology of overuse tendon injury in sports. Exhibit at American Academy of Orthopaedic Surgeons Annual Meeting, Washington, DC, February 20, 1992
86. Lin H, Ricci JL, Howard C, et al: Effects of growth factors on tendon cell growth. In *Trans Combined Meetings Orthop Res Soc USA, Japan and Canada*, October 21-23, 1991, p 12
87. Lundborg G, Rank F: Experimental intrinsic healing of flexor tendons based upon synovial fluid nutrition. *J Hand Surg* 3:21-31, 1978
88. Macierira-Coellto A, Tabury F: A re-evaluation of the changes in proliferations in human fibroblasts during ageing in vitro. *Cell Tissue Kinet* 15:213-224, 1982
89. Madden JW, Arem AJ: Wound healing: Biologic and clinical features. In Sabiston DC (ed): *Textbook of Surgery*, ed 14. Philadelphia, WB Saunders, 1991, p 167
90. Manske PR, Gelberman RH, Vande Berg JS, et al: Intrinsic flexor-tendon repair: A morphological study in vitro. *J Bone Joint Surg* 66A:385-396, 1984
91. Martens M, Wouters P, Burssens A, et al: Patellar tendinitis: Pathology and results of treatment. *Acta Orthop Scand* 53:445-450, 1982
92. Martinez-Hernandez A: Repair, degeneration, and fibrosis. In Rubin E, Faber JL (eds): *Pathology*. Philadelphia, JB Lippincott, 1988, pp 68-95
93. Mass DP, Tuel RJ, Labarbera M: Continuous oscillation of loaded and unloaded rabbit flexor tendons in vitro. In *Trans Combined Meetings Orthop Res Soc USA, Japan and Canada*, October 21-23, 1991, p 59
94. Mast BA, Krummel TM: Acute inflammation in fetal wound healing. In Adzick NS, Longaker MT (eds): *Fetal Wound Healing*. New York, Elsevier, 1992, pp 227-240
95. Menard D, Stanish WD: The ageing athlete. *Am J Sports Med* 17:187-196, 1989
96. Nagy IZ, Van Hahn HP, Verzar F: Age-related alterations in the cell nuclei and the DNA content of rat tail tendon. *Gerontologia* 15:258-264, 1969
97. Nemeth GG, Bolander ME, Martin GR: Growth factors and their role in wound and fracture healing. In *Growth Factors and Other Aspects of Wound Healing: Biological and Clinical Implications*. New York Alan R. Liss, 1988, pp 1-17
98. Neuberger A, Slack HGB: Metabolism of collagen from liver, bone, skin, and tendon in normal rat. *Biochem J* 53:47-52, 1953
99. Nirschl RP: Arm care: Relief and prevention for shoulder tendinitis. In *Tennis Elbow, Wrist Sprain in Athletes and Other Activities*. Arlington, Med Sports, 1981
100. Nirschl RP: Mesenchymal syndrome. *Virginia Medical Monthly* 96:659-662, 1969
101. Nirschl RP, Pettrone FA: Tennis elbow. The surgical treatment of lateral epicondylitis. *J Bone Joint Surg* 61A:832-839, 1979
102. Noyes FR: Functional properties of knee ligaments and alterations induced by

im
Or
103. Ne
(de
56
104. Oa
(St
105. Pe
Ch
106. Pe
fez
Tr
107. Pe
Ye
108. Pu
J S
109. Ra
Su
110. Ro
of
Pr
111. Ro
im
SL
Ri
112. Ro
su
113. Ro
bil
114. Ru
JB
115. Ru
116. Ru
Tr
117. Sc
Ar
118. Sc
119. Sc
mi
120. Si
He
CV
121. Sn
122. St
38
123. St
124. Sp
cli
125. Ta
co
19
126. Te
M
127. Tc
uli
128. Tr
14
129. UI
M

- immobilization: A correlative biomechanical and histological study in primates. *Clin Orthop* 123:210-242, 1977
103. Noyes FR, Lindenfeld TN, Marshall MT: What determines an athletic injury (definition)? Who determines an injury (occurrence)? *Am J Sports Med* 16(suppl): 565-566, 1988
 104. Oakes BW: Acute soft tissue injuries: Nature and management. *Aust Fam Physician (Suppl)* 10:3-16, 1982
 105. Perez-Tamayo R: Mechanisms of diseases: An Introduction to Pathology, ed 2, Chicago, Year Book Medical Publishers, 1985, p 182
 106. Perugia L, Ippolito E, Postacchini F: A new approach to the pathology. Clinical features and treatment of stress tendinopathy of the Achilles tendon. *Ital J Orthop Traumatol* 2:5-21, 1976
 107. Peterson L, Renstrom P: Sports Injuries. Their Prevention and Treatment. Chicago, Year Book Medical Publishers, 1986
 108. Puddu G, Ippolito E, Postacchini F: A classification of Achilles tendon disease. *Am J Sports Med* 4:145-150, 1976
 109. Rathbun JB, McNab I: The microvascular pattern of the rotator cuff. *J Bone Joint Surg* 52B:540-553, 1970
 110. Roberts AB, Sporn MB, et al: Transforming growth factor type B: Rapid induction of fibrosis and angiogenesis in vivo and stimulation of collagen formation in vitro. *Proc Natl Acad Sci USA* 83:4167-4171, 1986
 111. Rodosky MW, Fu FH: Induction of synovial inflammation by matrix molecules implant particles and chemical agents. In Leadbetter WB, Buckwalter JA, Gordon SL (eds): Sports-Induced Inflammation: Clinical and Basic Science Concepts. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1990, p 357
 112. Roughley PJ, White RJ: Age-related changes in the structure of the proteoglycan subunits from human articular cartilage. *J Biol Chem* 255:217-224, 1980
 113. Roy S, Irvin R: Sports Medicine: Prevention, Evaluation, Management, and Rehabilitation, Englewood Cliffs, NJ, Prentice Hall, 1983
 114. Rubin E, Faber JL: Cell injury. In Rubin E, Faber JL (eds): Pathology. Philadelphia, JB Lippincott, 1988, p 2
 115. Rubin E, Faber JL (eds): Pathology. Philadelphia, JB Lippincott, 1988
 116. Russell JE, Manske PR: Collagen synthesis during tendon repair: Epitenon impact. *Trans Orthop Res Soc* 14:274, 1989
 117. Scarpelli DG, Chiga M: Cell injury and errors of metabolism. In Kissahe JM (ed): Anderson's Pathology, vol 1, ed 8. St. Louis, CV Mosby, 1985, p 61
 118. Scarpelli DG, Trump BF: Cell Injury. Kalamazoo, MI, Upjohn Co, 1971
 119. Schatzker J, Branemark PI: Intravital observation on the microvascular anatomy and microcirculation of the tendon. *Acta Orthop Scand* 126(suppl):1-23, 1969
 120. Singer KM, Jones DC: Soft tissue conditions of the ankle and foot. In Nicholas JA, Herschman EB (eds): The Lower Extremity and Spine in Sports Medicine. St. Louis, CV Mosby, 1986, p 148
 121. Smith JW: Blood supply of tendons. *Am J Surg* 109:272-276, 1965
 122. Stanulis-Praeger BM: Cellular senescence revisited: A review. *Mech Ageing Dev* 38:1-48, 1987
 123. Stedman's Medical Dictionary, ed 25. Baltimore, Williams & Wilkins, 1990
 124. Sporn MB, Roberts AB: Transforming growth factor-B: Multiple actions and potential clinical applications. *JAMA* 262:938-941, 1989
 125. Takai S, Woo SL-Y, Gelberman RH, et al: The effect of frequency and duration on controlled passive mobilization of tendon healings. *Trans Orthop Res Soc* 16:137, 1991
 126. Teitz CC: Overuse injuries. In Teitz CC (ed): Scientific Foundation of Sports Medicine. Toronto, BC Decker, 1989, p 299
 127. Torp S, Baer E, Friedman B: Effects of age and mechanical deformation on the ultrastructures of tendon. Bristol, UK, University of Bristol, 1974, pp 223-249
 128. Trotter JA, Wofsy C: The length of collagen fibrils in tendon. *Trans Orthop Res Soc* 14:180, 1989
 129. Uthoff HK, Sarkar K: Classification and definition of tendinopathies. *Clin Sports Med* 10:707-720, 1991

130. van der Meulen JC: Present state of knowledge on processes of healing in collagen structures. *Int J Sports Med* 3(suppl 1):4-8, 1982
131. Virchow RLK: Cellular pathology as based upon physiological and pathological histology. New York, Donen, 1971
132. Vogel KG: Proteoglycans accumulate in a region of human tibialis posterior tendon subjected to compressive force in vitro and in ligaments. *In Trans Combined Meeting Orthop Res Soc USA, Japan and Canada, October 21-23, 1991, p 58*
133. Weinstein J: Neurogenic and non-neurogenic pains and inflammatory mediators. *Orthop Clin North Am* 22:235-246, 1991
134. Weiss JA, Woo SL-Y, Obland KJ, et al: Evaluation of a new injury model to study medical collateral ligament healing: Primary repair versus inoperative treatment. *J Orthop Res* 9:516-528, 1991
135. Weiss L: The cell. *In Weiss L (ed): Cell and Tissue Biology. A Textbook of Histology, ed 6. Baltimore, Urban and Schwarzenberg, 1988, p 22*
136. Williams JGP: Achilles tendon lesions in sports. *Sports Med* 3:114-135, 1986
137. Wilmore JH: Training for Sports and Activity: The Physiological Basis of the Conditioning Process, ed 2. Boston, Allyn & Bacon, 1982
138. Wilson AM, Goodship AE: Hysteresis energy losses in the equine superficial digital flexor tendon during exercise produce a local temperature sufficient to damage fibroblasts in vitro. *Trans Orthop Res Soc* 17:679, 1992
139. Woo SL-Y, Buckwalter JA (eds): Injury and Repair of Musculoskeletal Soft Tissues. Park Ridge, IL, American Academy of Orthopaedic Surgeons, 1988
140. Woo SL-Y, Matthews JV, Akeson WH, et al: Connective tissue response to immobility. Correlative study of biomechanical measurements of normal and immobilized rabbit knees. *Arthritis Rheum* 18:257-264, 1975
141. Zamora AJ, Marin JF: Tendon and myo-tendinous junction in an overloaded skeletal muscle of the rat. *Anat Embryol* 179:89-96, 1988

Address reprint requests to
Wayne B. Leadbetter, MD
The Orthopaedic Center
9711 Medical Center Drive, Suite 201
Rockville, MD 20850

TENDI

L
a mir
(CT)
prese:
Altho
US, C
strati
in th
asses:
the jo
cuff.

F
vast
princi
to sep
lutor
many
much
accur
tendc
clini:
respo
S
interp
graph
of an

From t

CLINIC

VOLUM

CLINICS IN SPORTS MEDICINE

Tendinitis I: Basic Concepts

PER A.F.H. RENSTRÖM, MD, PhD, and
WAYNE B. LEADBETTER, MD, GUEST EDITORS

VOLUME 11


• NUMBER 3 •

JULY 1992

W.B. SAUNDERS COMPANY
Harcourt Brace Jovanovich, Inc.

PHILADELPHIA LONDON TORONTO MONTREAL SYDNEY TOKYO

Accessible version


[Home](#) [About NINDS](#) [Disorders](#) [Funding & Research](#) [News & Events](#) [Find People](#) [Jobs & Training](#)
Science for the Brain

The nation's leading supporter of biomedical research on disorders of the brain and nervous system

More about Carpal Tunnel Syndrome

[Studies with patients](#)
[Research literature](#)
[Press releases](#)

DisclaimerSearch NINDS... [\(help\)](#)

Enter Words Here

[Go](#)**Contact Us****My Privacy**

NINDS is part of the
[National Institutes of Health](#)

You are here: [Home](#) > [Disorders](#) > [Carpal Tunnel Syndrome](#)**NINDS Carpal Tunnel Syndrome Information Page**Condensed from [Carpal Tunnel Syndrome Fact Sheet](#)[Get Web page suited for printing](#)[Email this to a friend or colleague](#)**Table of Contents (click to jump to sections)**[What is Carpal Tunnel Syndrome?](#)[Is there any treatment?](#)[What is the prognosis?](#)[What research is being done?](#)**Organizations**[Related NINDS Publications and Information](#)[Publicaciones en Español](#)[Additional resources from MEDLINEplus](#)**What is Carpal Tunnel Syndrome?**

Carpal tunnel syndrome occurs when tendons or ligaments in the wrist become enlarged, often from inflammation, after being aggravated. The narrowed tunnel of bones and ligaments in the wrist pinches the nerves that reach the fingers and the muscles at the base of the thumb. The first symptoms usually appear at night. Symptoms range from a burning, tingling numbness in the fingers, especially the thumb and the index and middle fingers, to difficulty gripping or making a fist, to dropping things. Some cases of carpal tunnel syndrome are due to work-related cumulative trauma of the wrist. Diseases or conditions that predispose to the development of carpal tunnel syndrome include pregnancy, diabetes, and obesity.

Is there any treatment?

Carpal tunnel syndrome is treated by immobilizing the wrist in a splint to minimize or prevent pressure on the nerves. If that fails, patients are sometimes given anti-inflammatory drugs or injections of cortisone in the wrist to reduce the swelling. There is also a surgical procedure in which doctors can open the wrist and cut the ligament at the bottom of the wrist to relieve the pressure. However, only a small percentage of patients require surgery.

What is the prognosis?

Approximately 1 percent of individuals with carpal tunnel syndrome develop permanent injury. The majority recover completely and can avoid reinjury by changing the way they do repetitive movements, the frequency with which they do the movements, and the amount of time they rest between periods when they perform the movements.

What research is being done?

Much of the on-going research on carpal tunnel syndrome is aimed at prevention and rehabilitation. The National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS) funds research on carpal tunnel syndrome.

[Select this link](#) to view a list of studies currently seeking patients.

Organizations

**American Academy of Orthopaedic Surgeons/
 American Association of Orthopaedic Surgeons**
 6300 North River Road
 Rosemont, IL 60018
hackett@aaos.org
<http://www.aaos.org>
 Tel: 847-823-7186
 Fax: 847-823-8125

**American Chronic Pain Association
 (ACPA)**
 P.O. Box 850
 Rocklin, CA 95677-0850
ACPA@pacbell.net
<http://www.theacpa.org>
 Tel: 916-632-0922 800-533-3231
 Fax: 916-632-3208

National Chronic Pain Outreach Association (NCPOA)
P.O. Box 274
Millboro, VA 24460
ncpoa@cfw.com
<http://www.chronicpain.org>
Tel: 540-862-9437
Fax: 540-862-9485

National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS)
National Institutes of Health, DHHS
31 Center Dr., Rm. 4C02 MSC 2350
Bethesda, MD 20892-2350
NIAMInfo@mail.nih.gov
<http://www.niams.nih.gov>
Tel: 301-496-8190 877-22-NIAMS (226-4267)

Centers for Disease Control and Prevention (CDCP)
U.S. Department of Health and Human Services
1600 Clifton Road, N.E.
Atlanta, GA 30333
inquiry@cdc.gov
<http://www.cdc.gov>
Tel: 800-311-3435 404-639-3311/404-639-3543

Occupational Safety & Health Administration
U.S. Department of Labor
200 Constitution Avenue, NW
Washington, DC 20210
<http://www.osha.gov>
Tel: 800-321-OSHA (-6742)

Related NINDS Publications and Information

- [Carpal Tunnel Syndrome Fact Sheet](#)
Carpal Tunnel Syndrome fact sheet compiled by the National Institute of Neurological Disorders and Stroke (NINDS).

Publicaciones en Español

- [Síndrome del Túnel Carpiano](#)
Información del Síndrome del Túnel Carpiano/Spanish-language fact sheet on carpal tunnel syndrome compiled by the National Institute of Neurological Disorders and Stroke (NINDS).

Prepared by:
Office of Communications and Public Liaison
National Institute of Neurological Disorders and Stroke
National Institutes of Health
Bethesda, MD 20892

NINDS health-related material is provided for information purposes only and does not necessarily represent endorsement by or an official position of the National Institute of Neurological Disorders and Stroke or any other Federal agency. Advice on the treatment or care of an individual patient should be obtained through consultation with a physician who has examined that patient or is familiar with that patient's medical history.

All NINDS-prepared information is in the public domain and may be freely copied. Credit to the NINDS or the NIH is appreciated.

[Return to top](#)

Last updated December 03, 2004

National Institute of Neurological Disorders and Stroke
[Home](#) | [About NINDS](#) | [Disorders](#) | [Funding](#) | [News and Events](#) | [Find People](#) | [Jobs and Training](#) | [Accessibility](#)



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.